

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C11D 17/00, A61K 7/50</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/39273</b> <b>(43) International Publication Date:</b> 06 July 2000 (06.07.2000)
<b>(21) International Application Number:</b> PCT/GB99/04328 <b>(22) International Filing Date:</b> 20 December1999 (20.12.1999) <b>(30) Priority Data:</b> 9828719.6 24 December1998 (24.12.1998) GB <b>(60) Parent Application or Grant</b> UNILEVER PLC [/]; (). UNILEVER NV [/]; (). HINDUSTAN LEVER LIMITED [/]; (). LYLE, Ian, Gardner [/]; (). SALMON, Tom, Matthew, Forrest [/]; (). SCHOPPER-MARTENS, Margrit [/]; (). WINTEROT, Elisabeth [/]; (). BAKER, Colin, J. ; ().	<b>Published</b>	
<b>(54) Title: DETERGENT COMPOSITION</b> <b>(54) Titre: COMPOSITION DETERGENTE</b>  <b>(57) Abstract</b> <p>A packaged aqueous self-foaming liquid cleansing composition comprising a base composition comprising a surfactant system comprising an anionic surfactant and an amphoteric or zwitterionic surfactant and a hydrophobic component, and a post-foaming agent, wherein the composition is substantially free of soap and thickens on the addition of the post foaming agent to the base composition.</p> <b>(57) Abrégé</b> <p>Cette invention concerne une composition de nettoyage aqueuse, auto-moussante, sous conditionnement. Cette composition comprend une composition de base renfermant un système tensioactif fait d'un tensioactif anionique, d'un tensioactif amphotérique ou zwitterionique et d'un composé hydrophobe, ainsi qu'un agent post-moussant. Cette composition est sensiblement exempte de savon et s'épaissit lorsque l'on ajoute l'agent post-moussant à la composition de base.</p>		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : C11D 17/00, A61K 7/50		A1	(11) International Publication Number: WO 00/39273
			(43) International Publication Date: 6 July 2000 (06.07.00)
(21) International Application Number: PCT/GB99/04328		(72) Inventors: LYLE, Ian, Gardner; Lever Faberge Deutschland GmbH, Hamburger Strasse 23, D-22083 Hamburg (DE). SALMON, Tom, Matthew, Forrest; Lever Faberge Deutschland GmbH, Alter Postweg 25, D-21614 Buxtehude (DE). SCHOPPER-MARTENS, Margrit; Lever Faberge Deutschland GmbH, Alter Postweg 25, D-21614 Buxtehude (DE). WINTEROT, Elisabeth; Lever Faberge Deutschland GmbH, Alter Postweg 25, D-21614 Buxtehude (DE).	
(22) International Filing Date: 20 December 1999 (20.12.99)			
(30) Priority Data: 9828719.6 24 December 1998 (24.12.98) GB			
(71) Applicant (for AE AU BB CA CY DM GB GD GH GM IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZA ZW only): UNILEVER PLC (GB/GB); Unilever House, Blackfriars, London EC4P 4BQ (GB).		(74) Agent: BAKER, Colin, J.; Eric Potter Clarkson, Park View House, 58 The Ropewalk, Nottingham NG1 5DD (GB).	
(71) Applicant (for all designated States except AE AU BB CA CY DM GB GD GH GM IE IL IN KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZA ZW): UNILEVER NV (NL/NL); Weena 455, NL-3013 AL Rotterdam (NL).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for IN only): HINDUSTAN LEVER LIMITED [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Mumbai 400 020, Maharashtra (IN).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: DETERGENT COMPOSITION			
(57) Abstract  A packaged aqueous self-foaming liquid cleansing composition comprising a base composition comprising a surfactant system comprising an anionic surfactant and an amphoteric or zwitterionic surfactant and a hydrophobic component, and a post-foaming agent, wherein the composition is substantially free of soap and thickens on the addition of the post foaming agent to the base composition.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Switzerland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Cote d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroun	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## Description

5

10

15

20

25

30

35

40

45

50

55

5

10

DETERGENT COMPOSITION

15

20

25

30

35

40

45

50

55

5 The present invention relates to detergent compositions  
suitable for topical application for cleansing the  
human body, such as the skin and hair. In particular,  
it relates to self foaming gel compositions,  
conveniently propellant driven and of the "bag-in-can"  
variety, which are suitable for use as cleansing  
compositions, and in particular for use as shower gels,  
which are formulated to give mild cleansing and  
optionally conditioning of the skin, and which are  
dispensed as a gel but generate a foam which is rich  
and feels luxurious to use.

Self foaming gel compositions formulated to cleanse the  
skin are well known. Such products may be considered  
superior to mousse type cleaning products, which suffer  
from the disadvantage that when applying a pre-formed  
mousse onto the skin, the mousse tends to break down  
and dissipate quickly, without providing substantive  
lather. In contrast, self-foaming compositions are  
dispensed initially as viscous liquids or gels, which  
generate copious amounts of lather as they are spread  
over the skin and optionally rubbed.

It is also known to formulate products which provide in  
particular a cleansing benefit, which are in the form  
of a bag in can gel composition. For example, WO  
97/03646 discloses a bag in can cleansing composition  
which contains a base composition which has a viscosity  
of at least 9,500 cps, which in practice is achieved by  
the use of a specified thickener. The composition is  
said to have superior properties by virtue of the  
relatively high viscosity, in terms of improved  
stability and the ability to incorporate higher levels  
of volatile self foaming agent.

5

10

15

20

25

30

35

40

45

50

55

A variety of problems have been found to be associated with such bag in can gel compositions. A particular problem relates to the manufacture of such compositions, and especially how to dose into a bag in can arrangement a gel composition which is not so viscous so as to cause problems in the dosing of that composition, and/or to require the use of specialist equipment to dose the composition. In particular, it is clear from the teaching of WO97/03646 that the composition is required to have a minimum viscosity of 9,500 cps. As such, the composition is difficult to process and dose on conventional filling lines.

An associated problem is that in preparing surfactant based gel compositions which subsequently have a self foaming volatile component (such as a liquifiable propellant gas) dosed into them, a number of these compositions experience a drop in viscosity when the self foaming volatile component is dosed into them. This can be disadvantageous, since it is desirable that the product as dispensed has a good, relatively viscous gel structure. This in turn is so that the dispensed product will not only have good manipulation properties for the user, but also so that the relatively rigid gel structure is able to accommodate a relatively high amount of self foaming agent (e.g. liquifiable propellant gas), and to entrap it relatively well. This is so that the resulting product foams well when lathered (by virtue of the relatively high content of volatile component), but also so that it does not foam prematurely after being dispensed.

It is thought that because of the phenomena described of thinning when the propellant gas is dosed that it is necessary for the base composition to have a relatively high initial viscosity, such that any thinning of the product when propellant gas is dosed which does occur

5

is not too deleterious to the properties of the final product as dispensed.

10

5

15

10

20

15

25

GB-A-2,213,160 describes a soap free post foaming gel composition comprising a major amount of water, 3-23% of a water soluble anionic alkali metal alkyl ether sulphate surfactant, 1-24% of dispersible nonionic ethoxylated fatty alcohol or fatty ester, and 5-20% saturated aliphatic hydrocarbon foaming agent, with the anionic surfactant and the ethoxylated fatty alcohol or ester being present in pre-determined ratios. Compositions according to this teaching are based on the specific combination of the anionic and nonionic surfactants, i.e. the alkyl ether sulphate and the nonionic ethoxylated fatty alcohol or fatty ester, and the criticality of each ingredient in the composition is clearly referred to.

30

20

35

25

40

30

A further problem which may be encountered relates to the stability of such cleansing compositions, especially those compositions which are based on non-soap surfactants. Regarding soap-based compositions, post foaming shaving gels employing soap based formulations are well known in the art. However, fundamentally it is desirable to create a cleansing composition based on non-soap surfactants, since by eliminating or reducing any soap to a low level the mildness of the composition may be improved. However, the inclusion of only a low level of soap means that it can be relatively difficult to structure and stabilise the cleansing composition.

45

35

50

We have found a way of formulating such self foaming gel or viscous liquid compositions such that they can deliver effective cleansing and optionally conditioning of the skin, are relatively easy to manufacture, and also have the desired stability and structure characteristics.

55



5

10

15

Thus, according to a first aspect of the invention, there is provided a packaged aqueous self foaming liquid cleansing composition comprising a base composition comprising a surfactant selected from anionic, nonionic, amphoteric, zwitterionic or cationic surfactants, and mixtures thereof, a hydrophobic component, and a post foaming agent, wherein the composition is substantially free of soap and thickens on the addition of the post foaming agent to the base composition.

20

25

25

Compositions according to the invention may conveniently be packaged in a so-called bag in can package, in which the cleansing composition is packaged inside a flexible bag, which is itself disposed in a can, with a suitable propellant gas located between the can and bag which pressurises the composition in the bag to a degree where it can be dispersed from suitable hardware.

30

35

40

Alternatively, the composition can be packaged in a pressurised bladder type package, in which the composition is packaged in what amounts to an elasticated bladder, which bladder maintains the contents of the bladder under positive pressure. This pressure is sufficient to cause the contents of the bladder to be dispensed via a closure mechanism located in the package.

30

45

35

50

In a highly preferred embodiment of the invention, the aqueous composition, shortly prior to being mixed with the self foaming agent, and subsequently being dosed into the product package, has a viscosity of less than 9,000 mPa.s, preferably less than 8,000 mPa.s, more preferably less than 5,000 mPa.s, and even more preferably less than 3,000 mPa.s. It is also highly preferred that packaged composition is prepared by

55

5

10

15

5

10

preparing the aqueous base portion of the composition as a somewhat viscous liquid, but excluding the post foaming agent. Just prior (say within 1 minute, preferably within 10 seconds, preferably within 1 second) to dosing the aqueous composition into the package of the arrangement (which in the case of a bag in can package is conveniently already located within the can), the post foaming agent is dosed into the aqueous composition, which is then rapidly dosed into a suitable container, such as e.g. a bag within the dispensing package.

20

15

25

However, compositions according to the invention typically thicken on the addition of self foaming agent to form a viscous gel, such that the dispensed composition has good handling and sensory properties, and also such that it retains the post foaming agent well after being dispensed.

30

35

40

45

20

25

30

35

50

By manufacturing the composition in this manner, it is possible to make the aqueous base such that it has a relatively low viscosity, and is therefore easy to process. However, introduction of the post foaming agent into the composition just prior to dosing the composition into the container ensures that any thickening of the composition, which is desirable bearing in mind that the form of the dispensed composition is as a viscous liquid or gel, does not occur until the composition is inside the container. This method therefore simplifies the manufacturing process. Gel compositions according to the invention are not simple propellant driven compositions, and may not be dispensed from conventional propellant driven aerosol packaging, since they are too viscous to dispense from such conventional single compartment propellant driven aerosol packaging. It is also important that the packaged composition does not have

55

5

any headspace, since if this were to occur it could cause the composition to be dispensed as a mousse.

10

5

15

10

20

25

20

30

35

25

40

30

45

35

50

55

Where the packaged composition utilises bag in can packaging, compositions according to the invention have two associated gases; a propellant gas and a post (self) foaming agent. The propellant gas is that which is contained within the can, but acts against the outside of the bag in which the composition is contained to dispense the composition when an actuator on the can is used. The propellant gas can be any suitable gas, but is conveniently a non-liquifiable propellant gas, such as compressed air, though any propellant which would function to dispense the composition would be suitable. The propellant gas is present in the packaged composition at any required and suitable level, but is typically present in the packaged product at levels sufficient to produce a satisfactory dispensing pressure, which will typically be 2-12 bar, more preferably 3-9 bar.

The other component required in such packaged compositions according to the invention is a post foaming agent. The post foaming agent is present in compositions according to the invention to allow the composition, which is dispensed in the form of a stable gel, once dispensed to generate a foam, as it evaporates on contact with a skin surface, thereby experiencing body heat. The generation of a foam provides a product which has various desirable consumer attributes, including ease of handling and spreading, and desirable sensory properties.

Suitable post foaming agents for inclusion in compositions according to the invention are volatile or liquifiable, and include (but are not limited to) hydrocarbons, such as isobutane and isopentane. Post foaming agents are present in packaged compositions

5 according to the invention at levels of 4-15%, more preferably 5-10% by weight of the packaged composition.

10 Suitable post foaming agents should be capable of being contained in compositions according to the invention as liquids, which may have been formed under  
5 the pressure to which the packaged composition has been subjected. As such, it is also desirable and may even be essential that packages in which the packaged  
15 composition is stored have no head space, to prevent the premature evaporation of the post foaming agent.  
10

20 Packaged compositions according to the invention contain a blend of hydrophobic components, including perfume oils and hydrophobic benefit agent components, and hydrocarbon post foaming agents, which are blended  
15 and matched, and chosen in conjunction with a suitable surfactant or blend of surfactants, in order to form a stable isotropic system. Compositions according to the invention are often microemulsions.  
25

30 The hydrophobic component in the composition may conveniently be a benefit agent, a perfume oil, or another hydrophobic component. The composition according to the invention may conveniently be suitable  
35 25 for cleansing, and in addition optionally moisturising, conditioning or protecting the skin. Where the hydrophobic component in the composition is a benefit agent, this may be included in the composition to moisturise, condition and/or protect the skin. By  
40 30 "benefit agent" is meant a substance that softens the skin (stratum corneum) and keeps it soft by retarding the decrease of its water content, and/or protects the skin. However, substantial deposition on the skin of a  
45 hydrophobic component or benefit agent is not necessarily a feature of compositions according to the  
35 invention.  
50

Preferred hydrophobic components which may be solid or liquid at room temperature, but in compositions according to the invention are to be found in liquid form (either by virtue of being liquid at room temperature themselves, or by being solubilised in a hydrophobic liquid component so as to provide a liquid composition) include:

- a) silicone oils, gums and modifications thereof such as linear and cyclic polydimethylsiloxanes, amino, alkyl alkylaryl and aryl silicone oils;
- b) fats and oils including natural fats and oils such as jojoba, soyabean, rice bran, avocado, almond, olive, sesame, persic, castor, coconut, mink oils; cacao fat, beef tallow, lard; hardened oils obtained by hydrogenating the aforementioned oils; and synthetic mono, di and triglycerides such as myristic acid glyceride and 2-ethylhexanoic acid glyceride;
- c) waxes such as carnauba, spermaceti, beeswax, lanolin and derivatives thereof;
- d) hydrophobic plant extracts;
- e) hydrocarbons such as liquid paraffins, petroleum jelly, microcrystalline wax, ceresin, squalene, squalane, and mineral oil;
- f) higher fatty acids such as lauric, myristic, palmitic, stearic, behenic, oleic, linoleic linolenic, lanolic, isostearic and polyunsaturated fatty acids (PUFA) acids;
- g) higher alcohols such as lauryl, cetyl, steryl, oleyl, behenyl, and 2-hexadecanol alcohol;
- h) esters such as cetyl octanoate, myristyl lactate, cetyl lactate, isopropyl myristate, myristyl myristate, isopropyl palmitate, isopropyl adipate, butyl stearate, decyl oleate, cholesterol isostearate, glycerol

5

10

15

20

25

30

35

40

45

35

50

55

monostearate, glycerol distearate, glycerol  
tristearate, alkyl lactate for example lauryl  
lactate, alkyl citrate and alkyl tartrate;

i) essential oils such as fish oils, mentha,  
jasmine, camphor, white cedar, bitter orange  
peel, ryu, turpentine, cinnamon, bergamont,  
citrus unshiu, calamus, pine, lavender, bay,  
clove, hiba, eucalyptus, lemon, starflower,  
thyme, peppermint, rose, sage, menthol,  
cineole, eugenol, citral, citronelle,  
borneol, linalool, geraniol, evening  
primrose, camphor, thymol, spirantol, pinene,  
limonene and terpenoid oils;

j) lipids such as cholesterol, ceramides,  
sucrose esters and pseudo-ceramides as  
described in European Patent Specification  
No. 556 957;

k) vitamins such as vitamin A and E, and vitamin  
alkyl esters, including vitamin C alkyl  
esters;

l) sunscreens such as octyl methoxyl cinnamate  
(Parsol MCX) and butyl methoxy benzoylmethane  
(Parsol 1789)

m) phospholipids;

n) perfume oils; and

o) mixtures of any of the foregoing components.

Particularly preferred hydrophobic benefit agents  
include esters such as isopropyl palmitate and  
myristate, and perfume oils.

The hydrophobic benefit agent is preferably present in  
an amount of from 0.1 to 20 wt%, more preferably 0.5 to  
10%, even more preferably 0.5 to 5 wt% by weight of the  
composition.

The surfactant or surfactants can be selected from any  
known surfactant suitable for topical application to

5

10

15

20

25

30

35

40

45


50

55

the human body, provided that they are blended in such a way as to form a stable isotropic system with the hydrophobic components of the composition. Mild surfactants, i.e. surfactants which do not damage the stratum corneum, the outer layer of skin, are particularly preferred. Because of their lathering properties, anionic surfactants are highly preferred components of compositions according to the invention.


Where the composition contains an anionic surfactant, it is preferable that the composition also contains a co-surfactant, which can be a nonionic, cationic amphoteric or zwitterionic surfactant. However, compositions according to the invention are substantially soap free; that is, they contain less than about 1% by weight of soap.

A preferred anionic surfactant is alkyl ether sulphate of formula:



where R is an alkyl group of 8 to 22 carbon atoms, n ranges from 0.5 to 10 especially from 1.5 to 8, and M is a solubilising cation.

Another preferred anionic surfactant is fatty acyl isethionate of formula:




where R is an alkyl or alkenyl group of 7 to 21 carbon atoms and M is a solubilising cation such as sodium, potassium, ammonium or substituted ammonium. Preferably at least three quarters of the RCO groups have 12 to 18 carbon atoms and may be derived from coconut, palm or a coconut/palm blend.


Other possible anionic surfactants include alkyl glyceryl ether sulphates, alkyl sulposuccinates, acyl taurates, acyl sarcosinates, alkyl sulphoacetates,

alkyl phosphates, alkyl phosphate esters, acyl lactylates, acyl glutamates and mixtures thereof.


Sulphosuccinates may be monoalkyl sulphosuccinates having the formula:




and amido-MEA sulphosuccinates of the formula:



Sarcosinates are generally indicated by the formula:



Taurates are generally identified by the formula:




Harsh surfactants such as primary alkane sulphonate or alkyl benzene sulphonate will generally be avoided.

If the surfactant comprises soap, the soap is preferably derived from materials with a  $C_{12}$  to  $C_{18}$  substantially saturated carbon chain and, preferably, is a potassium soap with a  $C_{12}$  to  $C_{18}$  carbon chain.

Suitable nonionic surfactants include alkyl polysaccharides, lactobionamides, ethyleneglycol esters, glycerol monoethers, polyhydroxyamides (glucamide), primary and secondary alcohol ethoxylates, especially the  $C_{8-20}$  aliphatic alcohols ethoxylated with an average of from 1 to 20 moles of ethylene oxide per mole of alcohol. Other suitable nonionics include oil in water emulsifiers such as PEG-40 hydrogenated castor oil.



Suitable further surfactant materials are zwitterionic detergents which have an alkyl or alkenyl group of 7 to 18 carbon atoms and comply with an overall structural formula:-



where R<sup>1</sup> is alkyl or alkenyl of 7 to 18 carbon atoms  
R<sup>2</sup> and R<sup>3</sup> are each independently alkyl, hydroxyalkyl or  
carboxyalkyl of 1 to 3 carbon atoms;


m is 2 to 4;

n is 0 or 1;


X is alkylene of 1 to 3 carbon atoms optionally substituted with hydroxyl; and

Y is  $-\text{CO}_2^-$  or  $-\text{SO}_3^-$ .

Zwitterionic surfactants within the above general formula include simple betaines of formula:-



and amido betaines of formula:




where  $m$  is 2 or 3.


In both formulae  $R^1$ ,  $R^2$  and  $R^3$  are as defined previously.

$R^1$  may, in particular, be a mixture of  $C_{12}$  and  $C_{14}$  alkyl groups derived from coconut so that at least half, preferably at least three quarters, of the group  $R^1$  has 10 to 14 carbon atoms.  $R^2$  and  $R^3$  are preferably methyl.

A further possibility is a sulphobetaine of formula:-




or



where  $m$  is 2 or 3, or variants of these in which

$-(CH_2)_3SO_3^-$  is replaced by



$R^1$ ,  $R^2$  and  $R^3$  in these formulae are as defined previously.

Mixtures of any of the foregoing surfactants may also be used.

The surfactant is preferably present at a level of from 10 to 35 wt%, preferably 12 to 30 wt% of the composition.

An optional component in compositions according to the invention is a cationic polymer, such as for example cationic celluloses.

Cationic cellulose is available from Amerchol Corp. (Edison, NJ, USA) in their Polymer JR (trade mark) and

5

10

15

LR (trade mark) series of polymers, as salts of hydroxyethyl cellulose reacted with trimethyl ammonium substituted epoxide, referred to in the industry (CTFA) as Polyquaternium 10. Another type of cationic cellulose includes the polymeric quaternary ammonium salts of hydroxyethyl cellulose reacted with lauryl dimethyl ammonium-substituted epoxide, referred to in the industry (CTFA) as Polyquaternium 24. These materials are available from Amerchol Corp. (Edison, NJ, USA) under the tradename Polymer LM-200.

20

25

A particularly suitable type of cationic polysaccharide polymer that can be used is a cationic guar gum derivative, such as guar hydroxypropyltrimonium chloride (Commercially available from Rhone-Poulenc in their JAGUAR trademark series).

30

35

Examples are JAGUAR C13S, which has a low degree of substitution of the cationic groups and high viscosity, JAGUAR C15, having a moderate degree of substitution and a low viscosity, JAGUAR C17 (high degree of substitution, high viscosity), JAGUAR C16, which is a hydroxypropylated cationic guar derivative containing a low level of substituent groups as well as cationic quaternary ammonium groups, and JAGUAR 162 which is a high transparency, medium viscosity guar having a low degree of substitution.

40

45

Preferably the cationic polymer is selected from cationic cellulose and cationic guar derivatives. Particularly preferred cationic polymers are JAGUAR C13S, JAGUAR C15, JAGUAR C17 and JAGUAR C16 and JAGUAR C162, especially Jaguar C13S.

50

Cationic polymer if present in compositions according to the invention is present at levels of 0.01-1.2%, more preferably 0.05-1.0%, even more preferably 0.1-0.5% by weight of the composition.

55

5

10

15

20

25

30

35

40

45

50

55

5 In some embodiments an auxiliary structurant or thickener may be added to the composition. Suitable structurant materials include swelling clays, for example laponite; cross-linked polyacrylates such as Carbopol (TM) (polymers available from Goodrich); acrylates and copolymers thereof; polyvinylpyrrolidone and copolymers thereof; polyethylene imines; polymeric carboxylates, consisting of and including modified and unmodified starches, unsubstituted guar gums, agars, alginates, xanthan gum, carrageenan, cellulose derivatives, exudate gum, gellan gum, gelatin, pectins and seed gums; gellants; and mixtures thereof.

15 Preferred thickeners for the composition include fumed silica; alkyl silicone waxes; aluminium silicate; fatty acids and derivatives thereof, in particular, fatty acid monoglyceride polyglycol ethers; polyammonium stearate; hydrotalcites; and mixtures thereof. A particularly preferred thickener is PEG-120 methyl glucose dioleate.

25 The composition may also comprise a viscosity modifying agent, ie a material which adjusts the viscosity of the composition to be that which is suitable for and preferred by consumers. Suitable materials include ethylene glycols, propylene glycols, salts such as sodium chloride and ammonium sulphate; and sucrose esters.

30 Further examples of structurants and thickeners are given in the International Cosmetic Ingredient Dictionary, Seventh Edition, 1997, published by CTFA (The Cosmetic, Toiletry & Fragrance Association), incorporated herein by reference.

50 Compositions of the invention may be formulated as propellant driven and dispensed products for washing

5

10

5

the skin, for example, bath or shower gels, hand washing compositions or facial washing liquids, and products for washing the hair, as well as post foaming shaving gels. Shower gels, and also post foaming shaving gels are particularly preferred product forms.

15

10

20

Other typical optional components of such compositions include opacifiers, preferably at 0.2 to 2.0 wt%; preservatives, preferably at 0.2 to 2.0 wt%; pH adjusters, typically at 0.05-2.0wt%; colourants, preferably at 0.05-2.0wt%; biological extracts, preferably at 0.05-2.0wt%, humectants, such as glycerol or sorbitol, preferably at 0.1-10.0wt%; and perfumes, preferably at 0.5 to 2.0 wt%.

15

25

The invention will be further illustrated by reference to the following non-limiting examples.

#### EXAMPLES

20

30

Examples 1-3 represent the aqueous phases of compositions according to the invention for packaging in a bag in can type of package.

35

40

45

50

55

INCI Name	Wt(%)		
	1	2	3
Sodium laureth sulphate, 2EO	10.0	12.5	10.0
Cocoamidopropyl betaine	4.0	5.0	4.0
Guar hydroxypropyltrimonium chloride	0.15	0.15	0.15
Citric acid	0.325	0.238	0.24
Sodium benzoate	0.5	0.5	0.5
Peach extract + Propylene glycol + Water	0.2	0.2	0.2
PEG-120 methyl glucose dioleate	1.75	1.68	1.68
Isopropyl palmitate	0.5	1.0	1.0
Sodium hydroxide	0.012	-	-
CI 47005 (dye)	0.0102	0.0102	0.0102
CI 45100 (dye)	0.00072	0.00072	0.0007
Perfume	1.0	1.0	1.0
PEG-40 Hydrog. castor oil	-	0.5	0.5
Water	to 100		
pH	4.96	5.02	4.92
Viscosity (mPA s)	4300	1100	80

Compositions 1-3 may be made according to the method described below.

5

In a suitable preparation method, for which the exact amounts of each component can be adjusted according to the amounts actually present in the final composition, 809g of water is heated to 50°C and 10g of sodium benzoate is first dissolved in this. To this is added 290g of sodium laureth sulphate 2EO and a further 522g of water, with stirring. 266g of cocoamidopropyl betaine is then added and stirring continued to dissolve the surfactants. 10g of isopropyl palmitate and 4g of peach extract are next mixed in. The colourants, dissolved in 10g water at 20°C, are then added. 3g of guar hydroxypropyltrimonium chloride and

5

10

5

36g of PEG-120 methyl glucose dioleate is dissolved in 20g of perfume oil, and added to the cooled mixture, with stirring. Finally the product pH is adjusted to 4.9 with citric acid (ca. 6.5g) and sufficient water is added to bring the total sample weight to 2kg (ca 13.5g).

15

10

20

15

25

20

30

#### EXAMPLES 4-9

35

25

Compositions 4-9 were prepared according to the preparation method described in conjunction with Examples 1-3 above, and were subjected to evaluation tests regarding their separation on addition of self foaming agent, turbidity and consistency.

40

30

The compositions were evaluated as described below.

#### Laboratory Evaluation Methods

45

35

Viscosities of the base formulations were measured at 30°C using a Haake VT-500 Viscotester, MV 2 Spindle Nos 3, 4, 5 depending on viscosity range. Shear rate was  $1s^{-1}$ .

50

55

5 The effects of mixing the various aqueous formulations  
4-9 with volatile foaming agents were evaluated by a  
simple laboratory test in which n-pentane was used as a  
10 model hydrocarbon liquid. 40g of the aqueous base was  
5 first weighed into a heavy glass screw-top bottle. To  
this was added 2.1 g or 4.4 g of n-pentane (to achieve  
15 5% or 10% volatile content by weight) and the bottle  
was immediately closed tightly and shaken vigorously  
for 1 minute. The sample was then left to stand  
10 overnight at room temperature, and was examined the  
following day. Separation, turbidity and consistency  
20 were evaluated on a 3-point scale



	4	5	6	7	8	9
SLES 2EO (70% ai)	14.5	18.125	14.5	18.125	18.125	14.5
Cocamidopropyl betaine (30% ai)	13.3	16.625	13.3	16.625	16.625	13.3
Guar hydroxypropyltrimonium chloride	0.15	0.15	0.15	0.15	0.15	0.15
Citric acid	0.325	0.325	0.35	0.325	0.325	0.325
Sodium benzoate	0.5	0.5	0.5	0.5	0.5	0.5
Sodium hydroxide	0.012	0.012	0.012			
Isopropyl palmitate	0.5	0.5	1	1.5	1	0.5
Peach extract	0.2	0.2	0.2	0.2	0.2	0.2
PEG-40 hydrog. Castor oil			0.5	0.5	0.5	
Perfume (Douceur 369) 1.0%, ex. IFP					1	
Perfume (Caresse 567) 1.0%, ex. IFP			1	1		
Perfume (Caresse 554) 1.0%, ex. IFP						1
Water	q.s.	q.s.	q.s.	q.s.	q.s.	q.s.
TOTAL	95	95	95	95	95	95

SUBSTITUTE SHEET (RULE 26)

5

10

15

5

10

20

15

25

30

35

40

45

50

55

Each of the base formulations 4-9, which was to provide 95% by weight of the aqueous base, was modified by the addition of a further sub composition (e.g. in the case of example 4, the addition of components a-h), in the amounts stated. The amounts of the components added in the sub compositions are stated as parts by weight of the aqueous base, with the balance to 100% by weight of the aqueous base being made up by the addition of extra water.

The compositions were then prepared as described above in relation to examples 1-3, dosing the composition with pentane, and assessing its stability, turbidity and consistency after 24 hours. The results are shown below. For dosage purposes, acceptable compositions had to have a base viscosity (prior to pentane dosage) of not more than 9,000 mPa.s, and preferably much less than this.

## Base-Formulation 4

	a	b	c	d	e	f	g	h
Isopropyl palmitate	0.50	0.80	0.50	0.50	0.80	1.00	1.00	0.90
PEG-120 methyl glucose diolate	1.80	2.50	2.00	2.00	2.00	2.50	2.00	2.50
Propylene Glycol	0.00	0.00	0.00	0.20	0.00	0.00	0.00	0.00

## Base Viscosity/mPas

Douceur 369	4640	795	5775	5400				
Mousseline 945	8150	1700	10000	8300				
Caresse 567	12700	3500	14450	14900	1370	853	133	1710

22

	sp/tur/co	sp/tur/co	sp/tur/co	sp/tur/co	sp/tur/co	sp/tur/co	sp/tur/co	sp/tur/co
Douceur 369								
Properties of Gel								
+ 5% Pentane	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+
+ 10% Pentane	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+
Mousseline 945								
Properties of Gel								
+ 5% Pentane	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+
+ 10% Pentane	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+
Caresse 567								
Properties of Gel								
+ 5% Pentane	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+
+ 10% Pentane	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+	-/-/+

sp = separation  
tur = turbidity  
co = consistency

-- = severe separation  
- = very turbid  
- = not thickened

- = slight separation  
- = slightly turbid  
+ = thickened

+ = no separation  
+ = clear  
++ = gelled

SUBSTITUTE SHEET (RULE 26)

5                   Suitable compositions after the stability tests  
described have no separation (ie +), are at least  
thickened or gelled (ie + or ++), and may be slightly  
10                   turbid or clear (ie - or +). The results indicate that  
5                   suitable compositions according to the criteria applied  
are compositions 4b (Douceur 369), 4d (Douceur 369), 4b  
(Mousseline 945) and 4h (Caresse 567), all with 5%  
15                   pentane.

## Base-Formulation 5

k	l	m	n	o	p
0.80	1.00	2.00	1.00	1.50	1.4
2.50	2.50	2.50	2.50	2.50	2.5
0.00	0.00	0.00	1.00	0.00	0.00

Isopropyl  
palmitate  
PEG-120 methyl  
glucose dioleate  
Propylene Glycol

Base Viscosity  
/mpas

39300	70	18550		2740
20600	81	18700	1300	
35000	170	26400	5750	

Douceur 369  
Mousseline 945  
Caresse 567

sp/tur/co	sp/tur/co	sp/tur/co	sp/tur/co	sp/tur/co	sp/tur/co
Douceur 369					
Properties of Gel					
+ 5% Pentane				+/-/+	
+10% Pentane				+/-/++	
Mousseline 945					
Properties of Gel					
+ 5% Pentane			+/-/+		
+10% Pentane			+/-/++		
Caresse 567					
Properties of Gel					
+ 5% Pentane			-/-/+		
+10% Pentane			--/--/++		

5

10

5

The results indicate that suitable compositions according to the invention under the criteria applied are compositions 5p (Douceur 369) and 5o (Mousseline 945) both with 5% pentane, and also 5p (Douceur 369) and 5o (Mousseline 945), both with 10% pentane.

15

20

25

30

35

40

45

50

55

5

26

10

15

20

25

30

35

40

45

50

55

Base-Formulation 6

	q	r	s
Isopropyl palmitate	1.00	1.00	1.00
PEG-120 methyl glucose dioleate	2.80	2.40	2.00
PEG-40 Hydrogenated castor oil	0.50	0.50	0.50
Base Viscosity /cPaa (Haake VT500 stage 5)	291	116	70

Base Formulation 7

	t	u	v	w
	2.00	2.00	1.50	1.50
	2.80	2.50	2.00	2.20
	0.50	0.50	0.50	0.50
	318	148	674	1130

Caresse 567  
Properties of  
Gel  
+ 5% Pentane  
+ 10% Pentane

+/-/+	+/-/+	+/-/+
+/-/+	+/-/+	+/-/+

+/-/+	+/-/+	+/-/+
+/-/+	+/-/+	+/-/+

5

10

5

The results indicate that suitable compositions according to the invention under the criteria applied are compositions 6q, 6r, 6s, 7t, 7u and 7w, all with Caresse 567 at 5% pentane, and compositions 6q, 7t, 7u, 7v and 7w, all with Caresse 567 at 10% pentane.

15

20

25

30

35

40

45

50

55



### Base-Formulation 8

	X	Y	Z
isopropyl	1.50	1.50	1.50
palmitate			
PEG-120 methyl	2.50	2.10	1.80
PEG-40			
hydrogenerated	0.50	0.50	0.50
castor oil			

Base Viscosity /mpas

Douceur 369	963	306	208
Mouseline 945			
Caresse 554			

	sp/tur/co	sp/tur/co	sp/tur/co
Douceur 369			
Properties of Gel			
+ 5% Pentane	+/-	-/+	-/+
+ 10% Pentane	-/-	+/-	+/-
Carress 554			
Properties of Gel			
+ 5% Pentane			
+ 10% Pentane			

The results indicate that suitable compositions according to the invention under the criteria applied are compositions 8y and 8z with Douceur 369 at 10% pentane, compositions 9Fa, 9Fb, 9Fc with Caresse 554 at 5% pentane, and composition 9Fc with Caresse 554 at 10% pentane.

As the skilled person will appreciate from the foregoing, compositions according to the invention may be adjusted by routine and minor adjustment of the hydrophobic component of the composition, such as the benefit agent, the self foaming agent and the perfume oil, to reliably produce satisfactory compositions.

It is also highly preferable that compositions according to the invention fulfil all of the separation, turbidity and consistency tests described above. The consistency test is significant, as the dispersed composition should be in the form of a gel or thick liquid. The nature of the package in which the composition is typically stored (eg bag in can, or elasticated bladder) suggests that the composition is not readily homogenised once packaged, for example by shaking so product separation is undesirable. The turbidity test is indicative of long term stability of the formulation.

#### EXAMPLES 10 - 12

The following compositions were prepared as outlined above:

	10	11	12
SLES.2EO	18.2	18.2	13.2
Coccoamidopropyl betaine	--	--	5.0
Arlatone T (PEG 40 sorbitan peroleate)	7.0	7.0	7.0
PEG-40 Hydrog. castor oil	--	--	--

PEG-120 methyl glucose dioleate	--	--	--
Isopropyl palmitate (Estol 1517)	--	--	--
Isopropyl myristate (Estol 1514)	3.0	3.0	3.0
Perfume (Caresse 567, ex. IFF)	2.0	2.0	2.0
Guar hydroxypropyltrimonium chloride	--	--	--
Citric acid	--	--	--
Sodium benzoate	--	--	--
Sorbitol	2.1	--	2.1
Water	To 100		

The compositions were dosed into bag in can packaging, and assessed for their properties.

In addition, as in examples 4-9 above, the compositions were assessed for nature of the gel they provided, when dosed with both 5% and 10% pentane as a post foaming agent. Classification of the gel is as outlined in conjunction with examples 4-9 above.

#### Results

Example	10	11	12
pH	6.9	7.0	6.4
Base viscosity (mPas)	128	122	251

5

10

Example		10	11	12
Separation/	5%	+ / + / + +	+ / + / + +	- - / - - / -
Turbidity/	pentane			
Consistency;	10%	+ / + / + +	+ / + / + +	
	pentane			

15

5

20

Examples 10 and 11 coincide with examples 11 and 12 of GB 2213160 referred to above. Example 12 coincides with example 11 of GB 2213160, but with 5.0% cocoamidopropyl betaine added. As can be seen, example 12 is unstable, indicating that amphoteric surfactants would not be expected to be usable in compositions according to the teaching of GB 2213160.

10

25

30

35

40

45

50

55

## Claims

5

10

15

20

25

30

35

40

45

50

55

5

CLAIMS

10

5

1. A packaged aqueous self foaming liquid cleansing composition comprising a base composition comprising a surfactant system comprising an anionic surfactant and an amphoteric or zwitterionic surfactant and a hydrophobic component, and a post foaming agent, wherein the composition is substantially free of soap and thickens on the addition of the post foaming agent to the base composition.

15

10

20

15

2. A packaged aqueous liquid cleansing composition according to claim 1, wherein the composition additionally comprises a nonionic or a cationic surfactant.

25

20

30

3. A packaged aqueous liquid cleansing composition according to claim 1 or claim 2, wherein the base composition has a viscosity prior to mixing with the self foaming agent and being dosed into the package of 9,000 mPa.s or less.

35

25

4. A packaged aqueous liquid cleansing composition according to claim 3, wherein the viscosity is less than 8,000 mPa.s.

40

30

5. A packaged aqueous liquid cleansing composition according to claim 4, wherein the viscosity is less than 3,000 mPa.s.

45

35

6. A packaged aqueous liquid cleansing composition according to Claims 1 to 5 wherein the composition is packaged in a bag in can container.

50

7. A packaged aqueous liquid cleansing composition according to Claim 6, wherein the post foaming agent is

55

5

dosed into the base composition within 10 seconds of the composition being dosed into the bag.

10

8. A packaged aqueous liquid cleansing composition according to claim 7, wherein the post foaming agent is dosed into the base composition within 1 second of the composition being dosed into the bag.

15

9. A packaged aqueous liquid cleansing composition according to any of Claims 6 to 8, additionally comprising a propellant gas in sufficient quantities to generate a dispensing pressure, which dispensing pressure is between 3 and 9 bar.

20

10. A packaged aqueous liquid cleansing composition according to Claims 1 to 5 wherein the composition is packaged in an elasticated bladder container.

25

11. A packaged aqueous liquid cleansing composition according to any of the preceding claims, wherein the post foaming agent is present in the composition at levels of 4 to 15% by weight of the composition.

30

12. A packaged aqueous liquid cleansing composition according to any of the preceding claims, wherein the composition is in the form of an isotropic blend.

35

13. A packaged aqueous liquid cleansing composition according to any of the preceding claims, wherein the composition is in the form of a microemulsion.

40

14. A packaged aqueous liquid cleansing composition according to any of the preceding claims, wherein the composition contains less than about 1% by weight of soap.

45

35

50

55

5 15. A packaged aqueous liquid cleansing composition  
according to any preceding claim, wherein the self  
foaming agent is a volatile or liquifiable hydrocarbon.

10 16. A packaged aqueous liquid cleansing composition  
5 according to any preceding claim, wherein there is no  
headspace in the packaged composition.

15 17. A packaged aqueous liquid cleansing composition  
10 according to any preceding claim, wherein the  
hydrophobic component comprises isopropyl palmitate or  
isopropyl myristate.

20 18. A packaged aqueous liquid cleansing composition  
15 according to any preceding claim, wherein the  
hydrophobic component comprises 0.1-20% by weight of  
the composition of a hydrophobic benefit agent.

25 19. A packaged aqueous liquid cleansing composition  
20 according to any preceding claim additionally  
comprising a cationic polymer.

30 20. A packaged aqueous liquid cleansing composition  
35 according to any preceding claim, wherein the  
composition is not separated, is thickened or gelled,  
and slightly turbid or clear after twenty four hours.

40 21. A packaged aqueous liquid cleansing composition  
30 according to Claim 20, wherein the composition is the  
base composition and the post foaming agent.

45 22. A packaged aqueous liquid cleansing composition  
35 according to Claim 21, wherein 40g of the base  
composition and 2.1g or 4.4g of the post foaming agent  
are dosed into a screw topped bottle, closed tightly  
and shaken vigorously for one minute before being left  
50 to stand at room temperature.



5

23. A packaged aqueous liquid cleansing composition according to Claim 22, wherein the post foaming agent is n-pentane.

10

15

20

25

30

35

40

45

50

55

# INTERNATIONAL SEARCH REPORT

Initial Application No  
PCT/GB 99/04328

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C11D17/00 A61K7/50

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C11D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 03646 A (CUSONS INT LTD ;HALL CHRISTOPHER JOHN (GB); YAQUB NAJEM (GB)) 6 February 1997 (1997-02-06) cited in the application page 3, paragraph 2 -page 8, paragraph 1	1,2,9,23
A	WO 96 09032 A (GILLETTE CO ;GEORGE ROBERT C (US); LASOTA ANDREW M (GB)) 28 March 1996 (1996-03-28) page 5, line 19 - line 36 page 7, line 21 -page 8, line 3 page 9, line 25 - line 32 page 12, line 17 - line 20 -/-	1-6,9, 15,23

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document relating to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

4 May 2000

Date of mailing of the international search report

15/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5518 Patendean 2  
NL - 2200 HV Rijswijk  
Tel: (+31-70) 540-2040, Tx: 31 651 epo nl,  
Fax: (+31-70) 540-3018

Authorized officer

Ketterer, M

Form PCT/ISA210 (second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT

Initial Application No  
PCT/GB 99/04328

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 38 39 349 A (COLGATE PALMOLIVE CO) 15 June 1989 (1989-06-15) page 3, line 12 - line 17 page 5, line 65 page 6, line 4 - line 9 claim 4 & GB 2 213 160 A cited in the application	1-7,10, 14,15,23
A	EP 0 259 843 A (JOHNSON & SON INC S C) 16 March 1988 (1988-03-16) page 3, line 39 - line 41; tables	1,23
A	EP 0 247 766 A (OSIPOW LLOYD I ;SPITZER JOSEPH G (US); MARRA DOROTHEA C (US)) 2 December 1987 (1987-12-02) examples	1,23
A	US 3 541 581 A (MONSON JAMES A) 17 November 1970 (1970-11-17) column 10, line 55 - line 69	1,3,23

# INTERNATIONAL SEARCH REPORT

information on patent family members

Int'l. Application No  
PCT/GB 99/04328

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9703646 A	06-02-1997	AU 6526696 A EP 0840596 A	18-02-1997 13-05-1998
WO 9609032 A	28-03-1996	US 5500211 A AT 174504 T AU 708643 B AU 3556195 A BR 9509074 A CA 2199829 A DE 69506725 D DE 69506725 T EP 0782436 A ES 2125658 T JP 10506110 T NZ 293184 A ZA 9507900 A	19-03-1996 15-01-1999 12-08-1999 09-04-1996 14-07-1998 28-03-1996 28-01-1999 22-07-1999 09-07-1997 01-03-1999 16-06-1998 26-08-1998 29-04-1996
DE 3839349 A	15-06-1989	US 4772427 A AT 399654 B AT 289988 A AU 2100288 A BE 1001545 A BR 8803939 A CA 1315637 A CH 678811 A DK 433988 A ES 2007997 A FI 883634 A, B FR 2623816 A GB 2213160 A, B GR 88100513 A, B IE 61585 B IT 1224719 B JP 1153624 A LU 87321 A MX 163643 B NL 8802031 A NO 173555 C PT 88252 A, B SE 8802800 A ZA 8805794 A	20-09-1988 26-06-1995 15-11-1994 01-06-1989 28-11-1989 13-03-1990 06-04-1993 15-11-1991 02-06-1989 01-07-1989 02-06-1989 02-06-1989 09-08-1989 31-10-1989 16-11-1994 18-10-1990 15-06-1989 08-03-1989 10-06-1992 03-07-1989 29-12-1993 10-11-1989 02-06-1989 25-04-1990
EP 0259843 A	16-03-1988	AU 600039 B AU 7961687 A BR 8707454 A DK 244488 A JP 1500669 T NO 881981 A NZ 221710 A PT 85661 A, B WO 8801876 A	02-08-1990 07-04-1988 13-09-1988 05-05-1988 09-03-1989 06-05-1988 26-02-1990 01-10-1987 24-03-1988
EP 0247766 A	02-12-1987	US 4726944 A DE 3785484 A	23-02-1988 27-05-1993
US 3541581 A	17-11-1970	GB 1279145 A NL 6912564 A	28-06-1972 22-02-1971